

Analysis of distributional Riemann curvature tensor in any dimension

Michael Neunteufel (TU Wien)

Jay Gopalakrishnan (Portland State University)

Joachim Schöberl (TU Wien)

Max Wardetzky (University of Göttingen)



International Workshop Vector- and Tensor-Valued Surface PDEs, Dresden, Nov 30th, 2023

Contents

Riemannian manifolds and Regge metric

Riemannian manifolds

Riemannian manifold (Ω, g) , $\Omega \subset \mathbb{R}^N$, g metric tensor

rect_metric.pdf

Riemannian manifolds

Riemannian manifold (Ω, g) , $\Omega \subset \mathbb{R}^N$, g metric tensor

Levi-Civita connection ∇

$$\nabla_X g(Y, Z) = g(\nabla_X Y, Z) + g(Y, \nabla_X Z)$$

rect_metric.pdf



Riemannian manifolds

Riemannian manifold (Ω, g) , $\Omega \subset \mathbb{R}^N$, g metric tensor

Levi-Civita connection ∇

$$\nabla_X g(Y, Z) = g(\nabla_X Y, Z) + g(Y, \nabla_X Z)$$

- Approximation of g on a triangulation
- Compute lengths and angles
- Regge calculus and finite elements



rect_metric.pdf

rect_appr_metric.pdf

rect_appr_metric_length

Riemannian manifolds




Riemannian manifold (Ω, g) , $\Omega \subset \mathbb{R}^N$, g metric tensor

Levi-Civita connection ∇

$$\nabla_X g(Y, Z) = g(\nabla_X Y, Z) + g(Y, \nabla_X Z)$$

- Approximation of g on a triangulation
- Compute lengths and angles
- Regge calculus and finite elements
- How to compute curvature? Convergence?

$$\|\mathcal{R}(g_h) - \mathcal{R}(g)\|_? \leq \mathcal{O}(h^?)$$



rect_metric.pdf

rect_appr_metric.pdf

rect_appr_metric_length.

Regge calculus and Regge metrics

non_flat_trigs.pdf

Regge calculus and Regge metrics

non_flat_trigs_red1.pdf

flat_trig_angle.pdf

Regge calculus and Regge metrics

non_flat_trigs_red2.pdf

nonflat_trig_angle.p



Regge calculus and Regge metrics

non_flat_trigs_red2.pdf

trig_edges.pdf

Regge calculus and Regge metrics



$$\text{Reg}_h^k = \{\varepsilon \in \mathcal{P}^k(\mathcal{T}, \mathbb{R}_{\text{sym}}^{d \times d}) \mid [\![t^\top \varepsilon t]\!]_E = 0 \text{ for all edges } E\}$$





$$\varphi_{E_i} = \nabla \lambda_j \odot \nabla \lambda_k, \quad t_j^\top \varphi_{E_i} t_j = c_i \delta_{ij}, \quad \varphi_{T_i} = \lambda_i \nabla \lambda_j \odot \nabla \lambda_k$$

Regge calculus and Regge metrics

$$\text{Reg}_h^k = \{\varepsilon \in \mathcal{P}^k(\mathcal{T}, \mathbb{R}_{\text{sym}}^{d \times d}) \mid [\![t^\top \varepsilon t]\!]_E = 0 \text{ for all edges } E\}$$



$$\varphi_{E_i} = \nabla \lambda_j \odot \nabla \lambda_k, \quad t_j^\top \varphi_{E_i} t_j = c_i \delta_{ij}, \quad \varphi_{T_i} = \lambda_i \nabla \lambda_j \odot \nabla \lambda_k$$



Definition distributional Riemann curvature tensor

Motivation Riemann curvature tensor I


Riemann curvature tensor:

$$\mathcal{R}(X, Y, Z, W) = g(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z, W)$$

$$\mathcal{R}_{ijkl} = \partial_i \Gamma_{jkl} - \partial_j \Gamma_{ikl} + \Gamma_{ik}^p \Gamma_{jpl} - \Gamma_{jk}^p \Gamma_{ipl}$$

Christoffel symbols: $\nabla_{\partial_j} \partial_k = \Gamma_{jk}^l \partial_l$, $\{\partial_i\}_{i=1}^N$ coordinate frame

$$\Gamma_{ij}^k(g) = g^{kl} \frac{1}{2} (\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij}) = g^{kl} \Gamma_{ijl}$$



Motivation Riemann curvature tensor I

Riemann curvature tensor:


$$\mathcal{R}(X, Y, Z, W) = g(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z, W)$$

$$\mathcal{R}_{ijkl} = \partial_i \Gamma_{jkl} - \partial_j \Gamma_{ikl} + \Gamma_{ik}^p \Gamma_{jp}{}^l - \Gamma_{jk}^p \Gamma_{ip}{}^l$$

Christoffel symbols: $\nabla_{\partial_j} \partial_k = \Gamma_{jk}^l \partial_l$, $\{\partial_i\}_{i=1}^N$ coordinate frame

$$\Gamma_{ij}^k(g) = g^{kl} \frac{1}{2} (\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij}) = g^{kl} \Gamma_{ijl}$$

Contribution: Element-wise curvature $\mathcal{R}_T := \mathcal{R}(g_h)|_T$ for $T \in \mathcal{T}$





Motivation Riemann curvature tensor II

Second fundamental form: F hyper-surface with g -normal vector $\hat{\nu}$


$$\mathbb{II}_{\hat{\nu}}(X, Y) = -g(\nabla_X \hat{\nu}, Y) = g(\hat{\nu}, \nabla_X Y), \quad X, Y \in \mathfrak{X}(F)$$

$$(\mathbb{II}_{\hat{\nu}})_{ij} = (\delta_i^l - \hat{\nu}_i \hat{\nu}^l) \Gamma_{lpk} \hat{\nu}^k (\delta_j^p - \hat{\nu}^p \hat{\nu}_j), \quad \hat{\nu}^i = \frac{1}{\|g^{-1}\nu\|} g^{ij} \nu_j$$

Metric g_h only tangential-tangential continuous $\Rightarrow \hat{\nu}_F^{T_+} \neq -\hat{\nu}_F^{T_-}$, $F = T_+ \cap T_-$



curved_triangles



curved_triangles_nonsmooth.pdf

Motivation Riemann curvature tensor II



Second fundamental form: F hyper-surface with g -normal vector $\hat{\nu}$

$$\mathbb{II}_{\hat{\nu}}(X, Y) = -g(\nabla_X \hat{\nu}, Y) = g(\hat{\nu}, \nabla_X Y), \quad X, Y \in \mathfrak{X}(F)$$


$$(\mathbb{II}_{\hat{\nu}})_{ij} = (\delta_i^l - \hat{\nu}_i \hat{\nu}^l) \Gamma_{lpk} \hat{\nu}^k (\delta_j^p - \hat{\nu}^p \hat{\nu}_j), \quad \hat{\nu}^i = \frac{1}{\|g^{-1}\nu\|} g^{ij} \nu_j$$

Metric g_h only tangential-tangential continuous $\Rightarrow \hat{\nu}_F^{T_+} \neq -\hat{\nu}_F^{T_-}$, $F = T_+ \cap T_-$

Contribution: Jump of second fundamental form $[\mathbb{II}]_F = \mathbb{II}_{\hat{\nu}_F^{T_+}} + \mathbb{II}_{\hat{\nu}_F^{T_-}}$ for $F \in \mathring{\mathcal{F}}$



curved_triangles.pdf



curved_triangles_nonsmooth.pdf

Motivation Riemann curvature tensor II

Second fundamental form: F hyper-surface with g -normal vector $\hat{\nu}$

$$\mathbb{II}_{\hat{\nu}}(X, Y) = -g(\nabla_X \hat{\nu}, Y) = g(\hat{\nu}, \nabla_X Y), \quad X, Y \in \mathfrak{X}(F)$$



$$(\mathbb{II}_{\hat{\nu}})_{ij} = (\delta_i^l - \hat{\nu}_i \hat{\nu}^l) \Gamma_{lpk} \hat{\nu}^k (\delta_j^p - \hat{\nu}^p \hat{\nu}_j), \quad \hat{\nu}^i = \frac{1}{\|g^{-1}\nu\|} g^{ij} \nu_j$$

Metric g_h only tangential-tangential continuous $\Rightarrow \hat{\nu}_F^{T_+} \neq -\hat{\nu}_F^{T_-}$, $F = T_+ \cap T_-$


Contribution: Jump of second fundamental form $[\mathbb{II}]_F = \mathbb{II}_{\hat{\nu}_F^{T_+}} + \mathbb{II}_{\hat{\nu}_F^{T_-}}$ for $F \in \mathring{\mathcal{F}}$

Motivation: Radial curvature equation

$$\mathcal{R}(X, \hat{\nu}, \hat{\nu}, Y) = (\nabla_{\hat{\nu}} \mathbb{II})(X, Y) - \mathbb{III}(X, Y), \quad X, Y \in \mathfrak{X}(F), \quad \mathbb{III}(X, Y) = \langle \nabla_X \hat{\nu}, \nabla_Y \hat{\nu} \rangle$$



curved_triangles.pptx




curved_triangles_nonsmooth.pptx

Motivation Riemann curvature tensor III

Angle defect:


At co-dimension 2 simplex E (Vertex in 2D, edge in 3D): 2-dimensional g -orthogonal plane

$$\Theta_E = 2\pi - \sum_{T \supset E} \arccos(g|_T(\hat{\mu}_E^{F_+}, \hat{\mu}_E^{F_-}))$$



Like classical angle defect for 2D manifolds

angle_defect_smooth.pdf




Motivation Riemann curvature tensor III

Angle defect:

At co-dimension 2 simplex E (Vertex in 2D, edge in 3D): 2-dimensional g -orthogonal plane


$$\Theta_E = 2\pi - \sum_{T \supset E} \arccos(g|_T(\hat{\mu}_E^{F_+}, \hat{\mu}_E^{F_-}))$$



Like classical angle defect for 2D manifolds

Contribution: Θ_E for $E \in \mathcal{E}$

angle_defect_smooth.pdf



Distributional (densitized) Riemann curvature tensor

Test space:

$$\mathcal{A}(\mathcal{T}) = \{A \in T_0^4(\mathcal{T}) \mid A(X, Y, Z, W) = -A(Y, X, Z, W) = -A(X, Y, W, Z) = A(Z, W, X, Y), \\ A(\cdot, \hat{\nu}, \hat{\nu}, \cdot) \text{ is single-valued for all } F \in \mathring{\mathcal{F}}, A(\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}) \text{ is single-valued for all } E \in \mathring{\mathcal{E}}\}$$

$$\mathring{\mathcal{A}}(\mathcal{T}) = \{A \in \mathcal{A}(\mathcal{T}) : A(\cdot, \hat{\nu}, \hat{\nu}, \cdot) \text{ vanishes on all } F \in \mathcal{F}_\partial\}$$

Distributional (densitized) Riemann curvature tensor

Test space:

$$\begin{aligned}\mathcal{A}(\mathcal{T}) = \{A \in T_0^4(\mathcal{T}) \mid A(X, Y, Z, W) &= -A(Y, X, Z, W) = -A(X, Y, W, Z) = A(Z, W, X, Y), \\ A(\cdot, \hat{\nu}, \hat{\nu}, \cdot) \text{ is single-valued for all } F \in \mathring{\mathcal{F}}, \quad &A(\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}) \text{ is single-valued for all } E \in \mathring{\mathcal{E}}\}\end{aligned}$$

$$\mathring{\mathcal{A}}(\mathcal{T}) = \{A \in \mathcal{A}(\mathcal{T}) : A(\cdot, \hat{\nu}, \hat{\nu}, \cdot) \text{ vanishes on all } F \in \mathcal{F}_\partial\}$$

Distributional densitized Riemann curvature tensor

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}_T, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle [\![\mathbb{I}]\!], A_{\cdot \hat{\nu} \hat{\nu} \cdot} \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\hat{\mu} \hat{\nu} \hat{\nu} \hat{\mu}} \omega_E, \quad A \in \mathring{\mathcal{A}}(\mathcal{T})$$

 GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of distributional Riemann curvature tensor in any dimension, *arXiv:2311.01603*.

Specialization to distributional Gauss curvature

Gauss curvature

$$K = \frac{\mathcal{R}(X, Y, Y, X)}{\|X\|_g \|Y\|_g - g(X, Y)^2} = \frac{\mathcal{R}_{1221}}{\det g}$$

Geodesic curvature

$$\kappa_{\hat{\nu}} = g(\hat{\nu}, \nabla_{\hat{\tau}} \hat{\tau}) = \mathbb{I}_{\hat{\nu}}(\hat{\tau}, \hat{\tau})$$

Define test function $A(X, Y, Z, W) = -v \omega(X, Y) \omega(Z, W)$, $v \in \mathring{\mathcal{V}} = \{u \in C^0(\Omega) \mid u|_{\partial\Omega} = 0\}$

Specialization to distributional Gauss curvature

Gauss curvature

$$K = \frac{\mathcal{R}(X, Y, Y, X)}{\|X\|_g \|Y\|_g - g(X, Y)^2} = \frac{\mathcal{R}_{1221}}{\det g}$$

Geodesic curvature

$$\kappa_{\hat{\nu}} = g(\hat{\nu}, \nabla_{\hat{\tau}} \hat{\tau}) = \mathbb{I}_{\hat{\nu}}(\hat{\tau}, \hat{\tau})$$

Define test function $A(X, Y, Z, W) = -v \omega(X, Y) \omega(Z, W)$, $v \in \mathring{\mathcal{V}} = \{u \in C^0(\Omega) \mid u|_{\partial\Omega} = 0\}$

Distributional densitized Gauss curvature

$$\widetilde{K\omega}(v) = \frac{1}{4} \widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T K_T v \omega_T + \sum_{F \in \mathcal{F}} \int_F [\kappa] v \omega_F + \sum_{E \in \mathring{\mathcal{E}}} \Theta_E v(E).$$

-  BERCHENKO-KOGAN, GAWLIK: Finite element approximation of the Levi-Civita connection and its curvature in two dimensions, *Found Comput Math* (2022).
-  GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of curvature approximations via covariant curl and incompatibility for Regge metrics, *SMAI J. Comput. Math.* (2023).

Specialization to distributional scalar curvature

Scalar curvature

$$S = g^{il} g^{jk} \mathcal{R}_{ijkl}$$

Mean curvature

$$H = \text{tr}(\mathbb{I}) = g^{ij} \mathbb{I}_{ij}$$

- Kulkarni-Nomizu product $\circledcirc : T_0^2(\Omega) \times T_0^2(\Omega) \rightarrow T_0^4(\Omega)$

$$(h \circledcirc k)(X, Y, Z, W) = h(X, W)k(Y, Z) + h(Y, Z)k(X, W) - h(X, Z)k(Y, W) - h(Y, W)k(X, Z)$$

- Define test function $A = v g \circledcirc g$, $v \in \mathring{\mathcal{V}}$

Specialization to distributional scalar curvature

Scalar curvature

$$S = g^{il} g^{jk} \mathcal{R}_{ijkl}$$

Mean curvature

$$H = \text{tr}(\mathbb{I}) = g^{ij} \mathbb{I}_{ij}$$

- Kulkarni-Nomizu product $\circledcirc : T_0^2(\Omega) \times T_0^2(\Omega) \rightarrow T_0^4(\Omega)$

$$(h \circledcirc k)(X, Y, Z, W) = h(X, W)k(Y, Z) + h(Y, Z)k(X, W) - h(X, Z)k(Y, W) - h(Y, W)k(X, Z)$$

- Define test function $A = v g \circledcirc g$, $v \in \mathring{\mathcal{V}}$

Distributional densitized scalar curvature

$$\widetilde{S}\omega(v) = \frac{1}{4} \widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T S_T v \omega_T + 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F [\![H]\!] v \omega_F + 2 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E v \omega_E$$



GAWLIK, N.: Finite element approximation of scalar curvature in arbitrary dimension,
arXiv:2301.02159.

Specialization to distributional Ricci curvature tensor

Ricci tensor: $\text{Ric}_{ij} = g^{ab}\mathcal{R}_{iabj}$

$A = g \oslash U$, $U \in \{V \in \mathcal{S}(\mathcal{T}) : V \text{ is } tt\text{- and } nn\text{-continuous}, V|_F \text{ and } V(\hat{\nu}, \hat{\nu}) \text{ vanish } \forall F \in \mathcal{F}_\partial\}$,

$$(g \oslash U)(X, \hat{\nu}, \hat{\nu}, Y) = U|_F(X, Y) + g|_F(X, Y)U(\hat{\nu}, \hat{\nu})$$

$$(g \oslash U)(\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}) = U(\hat{\mu}, \hat{\mu}) + U(\hat{\nu}, \hat{\nu}) = \text{tr}(U) - \text{tr}(U|_E).$$

Specialization to distributional Ricci curvature tensor

Ricci tensor: $\text{Ric}_{ij} = g^{ab}\mathcal{R}_{iabj}$

$A = g \oslash U$, $U \in \{V \in \mathcal{S}(\mathcal{T}) : V \text{ is } tt\text{- and } nn\text{-continuous}, V|_F \text{ and } V(\hat{\nu}, \hat{\nu}) \text{ vanish } \forall F \in \mathcal{F}_\partial\}$,

$$(g \oslash U)(X, \hat{\nu}, \hat{\nu}, Y) = U|_F(X, Y) + g|_F(X, Y)U(\hat{\nu}, \hat{\nu})$$

$$(g \oslash U)(\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}) = U(\hat{\mu}, \hat{\mu}) + U(\hat{\nu}, \hat{\nu}) = \text{tr}(U) - \text{tr}(U|_E).$$

Distributional densitized Ricci curvature tensor

$$\begin{aligned} \widetilde{\text{Ric}}\omega(U) &= \frac{1}{4}\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \text{Ric}_T, U \rangle \omega_T + \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle [\![\mathbb{II}]\!], U|_F + U(\hat{\nu}, \hat{\nu})g|_F \rangle \omega_F \\ &\quad + \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E(U(\hat{\nu}, \hat{\nu}) + U(\hat{\mu}, \hat{\mu})) \omega_E \end{aligned}$$

-  GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of distributional Riemann curvature tensor in any dimension, *arXiv:2311.01603*.

Error analysis

Integral representation of error

- **Goal:** Find integral representation of H^{-2} -error

parametrization $\tilde{g}(t) = g + t(g_h - g)$

$$\widetilde{\mathcal{R}\omega}(A)(g_h) - \widetilde{\mathcal{R}\omega}(A)(g) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega}(A)(\tilde{g}(t)) dt$$

Integral representation of error

- **Goal:** Find integral representation of H^{-2} -error

parametrization $\tilde{g}(t) = g + t(g_h - g)$

$$\widetilde{\mathcal{R}\omega}(A)(g_h) - \widetilde{\mathcal{R}\omega}(A)(g) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega}(A)(\tilde{g}(t)) dt$$

- **Problem:** test function $A = A_g$ depends on metric tensor

$$\begin{aligned} \mathcal{A}(\mathcal{T}) &= \{A \in T_0^4(\mathcal{T}) \mid A(X, Y, Z, W) = -A(Y, X, Z, W) = -A(X, Y, W, Z) = A(Z, W, X, Y), \\ &\quad A(\cdot, \hat{\nu}, \hat{\nu}, \cdot) \text{ is single-valued for all } F \in \mathring{\mathcal{F}}, \quad A(\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}) \text{ is single-valued for all } E \in \mathring{\mathcal{E}}\} \end{aligned}$$

Integral representation of error

- **Goal:** Find integral representation of H^{-2} -error

parametrization $\tilde{g}(t) = g + t(g_h - g)$

$$\widetilde{\mathcal{R}\omega}(A)(g_h) - \widetilde{\mathcal{R}\omega}(A)(g) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega}(A)(\tilde{g}(t)) dt$$

- **Problem:** test function $A = A_g$ depends on metric tensor

$$\begin{aligned} \mathcal{A}(\mathcal{T}) = \{A \in T_0^4(\mathcal{T}) \mid & A(X, Y, Z, W) = -A(Y, X, Z, W) = -A(X, Y, W, Z) = A(Z, W, X, Y), \\ & A(\cdot, \hat{\nu}, \hat{\nu}, \cdot) \text{ is single-valued for all } F \in \mathring{\mathcal{F}}, \quad A(\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}) \text{ is single-valued for all } E \in \mathring{\mathcal{E}}\} \end{aligned}$$

- **Solution:** Uhlenbeck trick

transform to g -independent test functions U with $A_g = \mathbb{A}_g(U)$

Uhlenbeck trick

$\mathcal{U}(\mathcal{T}) = \{ U \in \Gamma(\wedge^{N-2}(\mathcal{T}) \odot \wedge^{N-2}(\mathcal{T})) : U|_F(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single valued for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(F), F \in \mathring{\mathcal{F}},$
 $U|_E(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single valued for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(E), E \in \mathring{\mathcal{E}} \}$

$U \in \mathcal{U}(\mathcal{T})$ is **metric independent**

Uhlenbeck trick

$\mathcal{U}(\mathcal{T}) = \{ U \in \Gamma(\wedge^{N-2}(\mathcal{T}) \odot \wedge^{N-2}(\mathcal{T})) : U|_F(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single valued for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(F), F \in \mathring{\mathcal{F}},$
 $U|_E(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single valued for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(E), E \in \mathring{\mathcal{E}} \}$

$U \in \mathcal{U}(\mathcal{T})$ is **metric independent**

$$\mathbb{A} : \mathcal{U}(\mathcal{T}) \rightarrow T_0^4(\mathcal{T}), \quad \mathbb{A}(U)^{ijkl} = \hat{\varepsilon}^{\alpha_1 \dots \alpha_{N-2} ij} \hat{\varepsilon}^{\beta_1 \dots \beta_{N-2} kl} U_{\alpha_1 \dots \alpha_{N-2} \beta_1 \dots \beta_{N-2}}$$
$$\hat{\varepsilon}^{\alpha_1 \dots \alpha_{N-2} ij} = \frac{1}{\sqrt{\det g}} \varepsilon^{\alpha_1 \dots \alpha_{N-2} ij}, \quad \mathbb{A} = \mathbb{A}_g$$

Uhlenbeck trick

$\mathcal{U}(\mathcal{T}) = \{ U \in \Gamma(\wedge^{N-2}(\mathcal{T}) \odot \wedge^{N-2}(\mathcal{T})) : U|_F(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single valued for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(F), F \in \mathring{\mathcal{F}},$
 $U|_E(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single valued for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(E), E \in \mathring{\mathcal{E}} \}$

$U \in \mathcal{U}(\mathcal{T})$ is **metric independent**

$$\mathbb{A} : \mathcal{U}(\mathcal{T}) \rightarrow T_0^4(\mathcal{T}), \quad \mathbb{A}(U)^{ijkl} = \hat{\varepsilon}^{\alpha_1 \dots \alpha_{N-2} ij} \hat{\varepsilon}^{\beta_1 \dots \beta_{N-2} kl} U_{\alpha_1 \dots \alpha_{N-2} \beta_1 \dots \beta_{N-2}}$$
$$\hat{\varepsilon}^{\alpha_1 \dots \alpha_{N-2} ij} = \frac{1}{\sqrt{\det g}} \varepsilon^{\alpha_1 \dots \alpha_{N-2} ij}, \quad \mathbb{A} = \mathbb{A}_g$$

Lemma

The mapping \mathbb{A}_g is bijective and there holds

$$\mathcal{A}(\mathcal{T}) = \{ \mathbb{A}_g(U) : U \in \mathcal{U}(\mathcal{T}) \}.$$

Lemma

Let $\sigma := \dot{g}(t)$, $A \in \mathcal{A}(\mathcal{T})$, $(SA)(X, Y, Z, W) = A(X, Z, Y, W)$ swaps second with third argument. There holds

$$\begin{aligned}\dot{A}(X, Y, Z, W) = & -\text{tr}(\sigma)A(X, Y, Z, W) + A(\sigma(X, \cdot)^\sharp, Y, Z, W) + A(X, \sigma(Y, \cdot)^\sharp, Z, W) \\ & + A(X, Y, \sigma(Z, \cdot)^\sharp, W) + A(X, Y, Z, \sigma(W, \cdot)^\sharp),\end{aligned}$$

Evolution of distributional Riemann curvature

Lemma

Let $\sigma := \dot{g}(t)$, $A \in \mathcal{A}(\mathcal{T})$, $(SA)(X, Y, Z, W) = A(X, Z, Y, W)$ swaps second with third argument. There holds

$$\begin{aligned}\dot{A}(X, Y, Z, W) &= -\text{tr}(\sigma)A(X, Y, Z, W) + A(\sigma(X, \cdot)^\sharp, Y, Z, W) + A(X, \sigma(Y, \cdot)^\sharp, Z, W) \\ &\quad + A(X, Y, \sigma(Z, \cdot)^\sharp, W) + A(X, Y, Z, \sigma(W, \cdot)^\sharp),\end{aligned}$$

$$\frac{d}{dt}(\langle \mathcal{R}, A \rangle \omega_T)|_{t=0} = (2\langle \nabla^2 \sigma, S(A) \rangle + \langle \mathcal{R}(\sigma(\cdot, \cdot)^\sharp, \cdot, \cdot, \cdot), A \rangle - \frac{1}{2} \text{tr}(\sigma) \langle \mathcal{R}, A \rangle) \omega_T,$$

$$\frac{d}{dt}(\langle [\![\mathbb{I}]\!], A_{\cdot \hat{\nu} \hat{\nu} \cdot} \rangle \omega_F)|_{t=0} = \frac{1}{2} \langle [\![(\sigma(\hat{\nu}, \hat{\nu}) - \text{tr}(\sigma|_F))\mathbb{I} + 2(\nabla_F \sigma)(\hat{\nu}, \cdot)|_F - (\nabla_{\hat{\nu}} \sigma)|_F], A_{\cdot \hat{\nu} \hat{\nu} \cdot} \rangle \omega_F,$$

$$\frac{d}{dt}(\Theta_E A_{\hat{\mu} \hat{\nu} \hat{\nu} \hat{\mu}} \omega_E)|_{t=0} = -\frac{1}{2} \left(\sum_{F \supset E} [\![\sigma(\hat{\nu}, \hat{\mu})]\!]_F^E + \text{tr}(\sigma|_E) \Theta_E \right) A_{\hat{\mu} \hat{\nu} \hat{\nu} \hat{\mu}} \omega_E.$$

Evolution of distributional Riemann curvature

Proposition

Let $\sigma := \dot{g}$ and $A \in \mathring{\mathcal{A}}(\mathcal{T})$ with corresponding $U = \mathbb{A}^{-1}(A) \in \mathring{\mathcal{U}}(\mathcal{T})$. Then there holds

$$\frac{d}{dt} (\widetilde{\mathbb{A}^{-1}\mathcal{R}\omega})(g)(U)|_{t=0} = a_h(g; \sigma, U) + b_h(g; \sigma, U),$$

$$\begin{aligned} a_h(g; \sigma, U) &= \sum_{T \in \mathcal{T}} \int_T \left(\langle \mathcal{R}(\sigma(\cdot, \cdot)^\sharp, \cdot, \cdot, \cdot), \mathbb{A}(U) \rangle - \frac{1}{2} \text{tr}(\sigma) \langle \mathcal{R}, \mathbb{A}(U) \rangle \right) \omega_T \\ &\quad - 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F \left(\text{tr}(\sigma|_F) \langle [\![\mathbb{I}]\!], \mathbb{A}(U)_{\cdot \hat{\nu} \hat{\nu} \cdot} \rangle - [\![\mathbb{I}]\!] : \sigma|_F : \mathbb{A}(U)_{\cdot \hat{\nu} \hat{\nu} \cdot} \right) \omega_F \\ &\quad - 2 \sum_{E \in \mathring{\mathcal{E}}} \int_E \text{tr}(\sigma|_E) \Theta_E \mathbb{A}(U)_{\hat{\mu} \hat{\nu} \hat{\nu} \hat{\mu}} \omega_E \end{aligned}$$

$$[\![\mathbb{I}]\!] : \sigma|_F : \mathbb{A}(U)_{\cdot \hat{\nu} \hat{\nu} \cdot} = [\![\mathbb{I}]\!]_{ij} (\sigma|_F)^{jk} (\mathbb{A}(U)_{\cdot \hat{\nu} \hat{\nu} \cdot})_k^i.$$

Evolution of distributional Riemann curvature

Proposition

Let $\sigma := \dot{g}$ and $A \in \mathring{\mathcal{A}}(\mathcal{T})$ with corresponding $U = \mathbb{A}^{-1}(A) \in \mathring{\mathcal{U}}(\mathcal{T})$. Then there holds

$$\frac{d}{dt} (\widetilde{\mathbb{A}^{-1} \mathcal{R} \omega})(g)(U)|_{t=0} = a_h(g; \sigma, U) + b_h(g; \sigma, U),$$

$$\begin{aligned} b_h(g; \sigma, U) &= 2 \sum_{T \in \mathcal{T}} \int_T \langle \nabla^2 \sigma, S(\mathbb{A}(U)) \rangle \omega_T \\ &\quad + 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle [\![\sigma(\hat{\nu}, \hat{\nu}) \mathbb{I} + (\nabla_F \sigma)(\hat{\nu}, \cdot)|_F + \nabla_F(\sigma(\hat{\nu}, \cdot))|_F - (\nabla_{\hat{\nu}} \sigma)|_F], \mathbb{A}(U)_{\cdot \hat{\nu} \hat{\nu} \cdot}]\!] \omega_F \\ &\quad - 2 \sum_{E \in \mathring{\mathcal{E}}} \int_E \sum_{F \supset E} [\![\sigma(\hat{\nu}, \hat{\mu})]\!]_F^E \mathbb{A}(U)_{\hat{\mu} \hat{\nu} \hat{\nu} \hat{\mu}} \omega_E. \end{aligned}$$

$b_h(g; \sigma, U) = 2 \widetilde{\nabla^2 \sigma}(S \mathbb{A}(U))$ is the **distributional covariant incompatibility operator**
 $\text{inc}(\sigma)^{ij} = \text{curl}(\text{curl}(\sigma)^\top)^{ij} = \varepsilon^{ikl} \varepsilon^{jmn} \partial_k \partial_m \sigma_{ln}$

Integral representation

- **Goal:** Estimate $\|(\widetilde{\mathbb{A}^{-1}\mathcal{R}\omega})(g_h) - (\mathbb{A}^{-1}\mathcal{R}\omega)(g)\|_{H^{-2}}$
- **Integral representation:** $\tilde{g}(t) = g + t(g_h - g)$, $\sigma = \frac{d}{dt}\tilde{g}(t) = g_h - g$

$$((\widetilde{\mathbb{A}^{-1}\mathcal{R}\omega})(g_h) - (\mathbb{A}^{-1}\mathcal{R}\omega)(g))(U) = \int_0^1 a_h(\tilde{g}(t); \sigma, U) + b_h(\tilde{g}(t); \sigma, U) dt$$

- **Proof strategy idea:** Estimate integrand

$$|a_h(\tilde{g}(t); \sigma, U)| \lesssim \|\sigma\|_{L^2} \|U\|_{H^2} = \|g_h - g\|_{L^2} \|U\|_{H^2}$$

$$|b_h(\tilde{g}(t); \sigma, U)| \lesssim \|g_h - g\|_{L^2} \|U\|_{H^2}$$

Extract convergence rate: $\|g_h - g\| \lesssim h^{k+1}$

Distributional covariant incompatibility operator

Lemma

Let $\sigma \in \text{Reg}(\mathcal{T})$, $\Psi \in \mathcal{A}(\mathcal{T})$ a **smooth test function** with compact support, and g a **smooth metric tensor**. Then the **distributional covariant incompatibility** operator $\widetilde{\nabla^2\sigma}(S\Psi)$ is

$$\begin{aligned}\widetilde{\nabla^2\sigma}(S\Psi) = & \sum_{T \in \mathcal{T}} \left[\int_T \langle \nabla^2\sigma, S\Psi \rangle \omega_T + \int_{\partial T} \langle (\nabla_F \sigma)(\cdot, \hat{\nu}) + \nabla_F(\sigma(\hat{\nu}, \cdot)) - \nabla_{\hat{\nu}}\sigma \right. \\ & \quad \left. + \sigma(\hat{\nu}, \hat{\nu}) \mathbb{I}_{\hat{\nu}}, (S\Psi)_{\cdot, \hat{\nu}, \hat{\nu}, \cdot} \rangle \omega_{\partial T} \right] - \sum_{E \in \mathring{\mathcal{E}}} \sum_{F \supset E} \int_E [\![\sigma(\hat{\nu}, \hat{\mu})]\!]_F^E (S\Psi)_{\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}} \omega_E.\end{aligned}$$

Distributional covariant incompatibility operator

Lemma

Let $\sigma \in \text{Reg}(\mathcal{T})$, $\Psi \in \mathcal{A}(\mathcal{T})$ a **smooth test function** with compact support, and g a **smooth metric tensor**. Then the **distributional covariant incompatibility** operator $\widetilde{\nabla^2\sigma}(S\Psi)$ is

$$\begin{aligned}\widetilde{\nabla^2\sigma}(S\Psi) = & \sum_{T \in \mathcal{T}} \left[\int_T \langle \nabla^2\sigma, S\Psi \rangle \omega_T + \int_{\partial T} \langle (\nabla_F \sigma)(\cdot, \hat{\nu}) + \nabla_F(\sigma(\hat{\nu}, \cdot)) - \nabla_{\hat{\nu}}\sigma \right. \\ & \quad \left. + \sigma(\hat{\nu}, \hat{\nu}) \mathbb{I}_{\hat{\nu}}, (S\Psi)_{\cdot, \hat{\nu}, \hat{\nu}, \cdot} \rangle \omega_{\partial T} \right] - \sum_{E \in \mathring{\mathcal{E}}} \sum_{F \supset E} \int_E [\![\sigma(\hat{\nu}, \hat{\mu})]\!]_F^E (S\Psi)_{\hat{\mu}, \hat{\nu}, \hat{\nu}, \hat{\mu}} \omega_E.\end{aligned}$$

Distributional covariant incompatibility operator

Lemma

Let $\sigma \in \text{Reg}(\mathcal{T})$, $\Psi \in \mathcal{A}(\mathcal{T})$ a **smooth test function** with compact support, and g a **smooth metric tensor**. Then the **distributional covariant incompatibility** operator $\widetilde{\nabla^2\sigma}(S\Psi)$ is

$$\begin{aligned}\widetilde{\nabla^2\sigma}(S\Psi) = & \sum_{T \in \mathcal{T}} \left[\int_T \langle \nabla^2\sigma, S\Psi \rangle \omega_T + \int_{\partial T} \langle (\nabla_F \sigma)(\cdot, \hat{\nu}) + \nabla_F(\sigma(\hat{\nu}, \cdot)) - \nabla_{\hat{\nu}}\sigma \right. \\ & \quad \left. + \sigma(\hat{\nu}, \hat{\nu}) \mathbb{I}_{\hat{\nu}}, (S\Psi)_{\cdot \hat{\nu} \hat{\nu} \cdot} \rangle \omega_{\partial T} \right] - \sum_{E \in \mathring{\mathcal{E}}} \sum_{F \supset E} \int_E [\sigma(\hat{\nu}, \hat{\mu})]_F^E (S\Psi)_{\hat{\mu} \hat{\nu} \hat{\nu} \hat{\mu}} \omega_E.\end{aligned}$$

Definition (incompatibility operator)

Let U such that $U = \mathbb{A}^{-1}(A)$ with $A \in \mathcal{A}(\Omega)$. For a symmetric matrix $\sigma \in T_0^2(\Omega)$ we define the **covariant incompatibility operator** $\text{inc } \sigma$ by

$$\langle \text{inc } \sigma, U \rangle = -\langle \nabla^2\sigma, S(A) \rangle, \quad \text{for all } A \in \mathcal{A}(\Omega).$$

Adjoint of distributional covariant incompatibility operator

Motivation:

$$|b_h(\tilde{g}(t); \sigma, U)| = \left| 2 \widetilde{\nabla^2 \sigma}((S\mathbb{A})(U)) \right| \lesssim \|\sigma\|_{L^2} \|U\|_{H^2}$$

Adjoint of distributional covariant incompatibility operator

Motivation:

$$|b_h(\tilde{g}(t); \sigma, U)| = \left| 2 \widetilde{\nabla^2 \sigma}((S\mathbb{A})(U)) \right| = \left| 2 \left(\text{divdiv} \widetilde{(S\mathbb{A})}(U) \right)(\sigma) \right| \lesssim \|\sigma\|_{L^2} \|U\|_{H^2}$$

Lemma

Let $\sigma \in \text{Reg}(\mathcal{T})$, $A \in \mathring{\mathcal{A}}(\mathcal{T})$, and g a Regge metric. There holds $\widetilde{\nabla^2 \sigma}(SA) = \widetilde{\text{divdiv}}(SA)(\sigma)$ with

$$\begin{aligned} \widetilde{\text{divdiv}}(SA)(\sigma) &= \sum_{T \in \mathcal{T}} \left[\int_T \langle \sigma, \text{divdiv}(SA) \rangle \omega_T + \int_{\partial T} (\langle \sigma|_F, (\text{div}(SA) + \text{div}_F(SA))_{\hat{\nu}} + H(SA)_{\hat{\nu}\hat{\nu}}) \right. \\ &\quad \left. - \sigma|_F : \mathbb{I} : (SA)_{\hat{\nu}\hat{\nu}} - \langle \mathbb{I} \otimes \sigma|_F, SA \rangle \right] \omega_{\partial T} - \sum_{E \in \mathring{\mathcal{E}}} \sum_{F \supset E} \int_E \langle \sigma|_E, \llbracket (SA)_{\hat{\nu}\hat{\mu}} \rrbracket_F^E \rangle \omega_E. \end{aligned}$$

Analysis of a_h and b_h

Proposition

Let $\tilde{g}(t) = g + (g_h - g)t$, $\sigma = g_h - g$, and $U \in H_0^2(\Omega, \mathcal{U})$. There holds for all $t \in [0, 1]$

$$|a_h(\tilde{g}(t); \sigma, U)| \lesssim (1 + \max_{T \in \mathcal{T}_h} h_T^{-1} \|g_h - g\|_{W^{1,\infty}(T)} + \max_{T \in \mathcal{T}_h} h_T^{-2} \|g_h - g\|_{L^\infty(T)}) \|\sigma\|_2 \|U\|_{H^2}.$$

Assume that $g_h = \mathcal{I}_h^k g$ is an optimal-order interpolant. Then for an integer $k \geq 1$

$$|a_h(\tilde{g}(t); \sigma, U)| \lesssim \left(\sum_{T \in \mathcal{T}_h} h_T^{p(k+1)} |g|_{W^{k+1,p}(T)}^p \right)^{1/p} \|U\|_{H^2} \approx h^{k+1} |g|_{W^{k+1,p}} \|U\|_{H^2}.$$

$$\|\sigma\|_2^2 = \|\sigma\|_{L^2}^2 + h^2 \|\sigma\|_{H_h^1}^2 + h^4 \|\sigma\|_{H_h^2}^2, \quad \|\sigma\|_{H_h^1}^2 = \sum_{T \in \mathcal{T}} \|\sigma\|_{H^1(T)}^2$$

Analysis of a_h and b_h

Proposition

Let $\tilde{g}(t) = g + (g_h - g)t$, $\sigma = g_h - g$, and $U \in H_0^2(\Omega, \mathcal{U})$. There holds for all $t \in [0, 1]$ for dimension $N \geq 3$

$$|b_h(\tilde{g}(t); \sigma, U)| \lesssim (1 + \max_{T \in \mathcal{T}_h} h_T^{-2} \|g_h - g\|_{L^\infty(T)} + \max_{T \in \mathcal{T}_h} h_T^{-1} \|g_h - g\|_{W^{1,\infty}(T)}) \|g_h - g\|_2 \|U\|_{H^2}$$

and for $N = 2$

$$|b_h(\tilde{g}(t); \sigma, U)| \lesssim (1 + \max_{T \in \mathcal{T}_h} h_T^{-1} \|g_h - g\|_{L^\infty(T)} + \|g_h - g\|_{W_h^{1,\infty}}) \|g_h - g\|_2 \|U\|_{H^2}.$$

Assume that $g_h = \mathcal{I}_h^k g$ is an optimal-order interpolant. Then for an integer $k \geq 1$ for $N \geq 3$ and $k \geq 0$ for $N = 2$

$$|b_h(\tilde{g}(t); \sigma, U)| \lesssim \left(\sum_{T \in \mathcal{T}_h} h_T^{p(k+1)} |g|_{W^{k+1,p}(T)}^p \right)^{1/p} \|U\|_{H^2} \approx h^{k+1} |g|_{W^{k+1,p}} \|U\|_{H^2}.$$

Results

Theorem (Gopalakrishnan, N., Schöberl, Wardetzky 2023)

Assume $\{g_h\}_{h>0}$ is a family of Regge metrics on a shape regular family of triangulations $\{\mathcal{T}_h\}_{h>0}$ with $\lim_{h \rightarrow 0} \|g_h - g\|_{L^\infty} = 0$ and $\sup_{h>0} \max_{T \in \mathcal{T}_h} \|g_h\|_{W^{2,\infty}(T)} < \infty$. Then there exists $h_0 > 0$ such that for all $h \leq h_0$ in the two-dimensional case $N = 2$

$$\|(\widetilde{\mathbb{A}^{-1}\mathcal{R}\omega})(g_h) - (\mathbb{A}^{-1}\mathcal{R}\omega)(g)\|_{H^{-2}} \lesssim \left(1 + \max_{T \in \mathcal{T}_h} (h_T^{-1} \|g - g_h\|_{L^\infty(T)}) + \|g - g_h\|_{W_h^{1,\infty}}\right) \|g_h - g\|_2$$

and for higher dimensions $N \geq 3$

$$\begin{aligned} & \|(\widetilde{\mathbb{A}^{-1}\mathcal{R}\omega})(g_h) - (\mathbb{A}^{-1}\mathcal{R}\omega)(g)\|_{H^{-2}} \\ & \lesssim \left(1 + \max_{T \in \mathcal{T}_h} (h_T^{-2} \|g - g_h\|_{L^\infty(T)}) + \max_{T \in \mathcal{T}_h} (h_T^{-1} \|g - g_h\|_{W^{1,\infty}(T)})\right) \|g_h - g\|_2. \end{aligned}$$

$$a_h(g; \sigma, U) = 0 \text{ for } N = 2$$

Results

Theorem (Gopalakrishnan, N., Schöberl, Wardetzky 2023)

Let k be an integer with $k \geq 0$ for $N = 2$ and $k \geq 1$ for $N \geq 3$. Assume that

$g_h = \mathcal{I}_h^k g \in \text{Reg}_h^k$ is a family of optimal order interpolants on a shape regular family of triangulations $\{\mathcal{T}_h\}_{h>0}$ with $\sup_{h>0} \max_{T \in \mathcal{T}_h} \|g_h\|_{W^{2,\infty}(T)} < \infty$. Then there exists $h_0 > 0$ such that for all $h \leq h_0$ and $p \in [2, \infty]$ satisfying $p > \frac{m}{k+1}$

$$\|(\widetilde{\mathbb{A}^{-1}\mathcal{R}\omega})(g_h) - (\mathbb{A}^{-1}\mathcal{R}\omega)(g)\|_{H^{-2}} \lesssim \left(\sum_{T \in \mathcal{T}_h} h_T^{p(k+1)} |g|_{W^{k+1,p}(T)}^p \right)^{1/p} \approx h^{k+1} |g|_{W^{k+1,p}}$$

where m is the codimension index of \mathcal{I}_h^k .

$$a_h(g; \sigma, U) = 0 \text{ for } N = 2$$

Specialization to 2D

Lemma

For $N = 2$ the distributional densitized Riemann curvature tensor simplifies to the distributional Gauss curvature

$$\widetilde{K\omega}(u) = \sum_{T \in \mathcal{T}} \int_T K_T u \omega_T + \sum_{F \in \mathring{\mathcal{F}}} \int_F [\kappa]_F u \omega_F + \sum_{E \in \mathring{\mathcal{E}}} \Theta_E u(E), \quad u \in \mathring{\mathcal{V}},$$

and there holds $\mathcal{U}(\mathcal{T}) = \mathring{\mathcal{V}}$ and

$$a_h(g; \sigma, u) = 0,$$

$$\begin{aligned} b_h(g; \sigma, u) = & -2 \sum_{T \in \mathcal{T}} \int_T \text{inc } \sigma u \omega_T + 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F [\text{curl}(\sigma)(\hat{\tau}) + \nabla_{\hat{\tau}}(\sigma(\hat{\nu}, \hat{\tau}))]_F u \omega_F \\ & - 2 \sum_{E \in \mathring{\mathcal{E}}} \sum_{F \supset E} [\sigma(\hat{\nu}, \hat{\mu})]_F^E u(E). \end{aligned}$$

Specialization to 3D

Curvature operator

$$\mathcal{Q}(X \wedge Y, W \wedge Z) := \mathcal{R}(X, Y, Z, W), \quad X \wedge Y \in \Lambda^2(\Omega)$$

$$\tilde{\mathcal{Q}}(\cdot, \cdot) := \mathcal{Q}(\star \cdot, \star \cdot) \in T_2^0(\Omega), \quad \mathcal{U}(\mathcal{T}) = \text{Reg}(\mathcal{T})$$

Lemma

$$\widetilde{\tilde{\mathcal{Q}}\omega}(U) = \sum_{T \in \mathcal{T}} \int_T \langle \tilde{\mathcal{Q}}_T, U \rangle \omega_T + \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle [\![\mathbb{I}]\!], (\hat{\nu} \otimes \hat{\nu}) \times U \rangle \omega_F + \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E U(\hat{\tau}, \hat{\tau}) \omega_E$$

$$\begin{aligned} a_h(g; \sigma, U) &= -2 \sum_{T \in \mathcal{T}} \int_T \tilde{\mathcal{Q}} : \sigma : U \omega_T - 2 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E \sigma(\hat{\tau}, \hat{\tau}) U(\hat{\tau}, \hat{\tau}) \omega_E \\ &\quad - 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F \left(\text{tr}(\sigma|_F) \langle [\![\mathbb{I}]\!], (\hat{\nu} \otimes \hat{\nu}) \times U \rangle - [\![\mathbb{I}]\!] : \sigma|_F : ((\hat{\nu} \otimes \hat{\nu}) \times U) \right) \omega_F \end{aligned}$$

Specialization to 3D

Curvature operator

$$\mathcal{Q}(X \wedge Y, W \wedge Z) := \mathcal{R}(X, Y, Z, W), \quad X \wedge Y \in \Lambda^2(\Omega)$$

$$\tilde{\mathcal{Q}}(\cdot, \cdot) := \mathcal{Q}(\star \cdot, \star \cdot) \in T_2^0(\Omega), \quad \mathcal{U}(\mathcal{T}) = \text{Reg}(\mathcal{T})$$

Lemma

$$\widetilde{\tilde{\mathcal{Q}}\omega}(U) = \sum_{T \in \mathcal{T}} \int_T \langle \tilde{\mathcal{Q}}_T, U \rangle \omega_T + \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle [\![\mathbb{I}]\!], (\hat{\nu} \otimes \hat{\nu}) \times U \rangle \omega_F + \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E U(\hat{\tau}, \hat{\tau}) \omega_E$$

$$\begin{aligned} b_h(g; \sigma, U) &= -2 \sum_{T \in \mathcal{T}} \int_T \langle \text{inc } \sigma, U \rangle \omega_T - 2 \sum_{E \in \mathring{\mathcal{E}}} \int_E \sum_{F \supset E} \llbracket \sigma(\hat{\nu}, \hat{\mu}) \rrbracket_F^E U(\hat{\tau}, \hat{\tau}) \omega_E \\ &\quad + 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket (\sigma(\hat{\nu}, \hat{\nu}) \mathbb{I} + \nabla_F(\sigma(\hat{\nu}, \cdot))) \times (\nu \otimes \nu) + Q(\text{curl } \sigma)^\top \times \hat{\nu} \rrbracket, U|_F \rangle \omega_F \end{aligned}$$

Numerical examples

3D curvature

$$\Omega = (-1, 1)^3$$

$$\Phi(x, y, z) = (x, y, z, f(x, y, z)), \quad f(x, y, z) = \frac{1}{2}(x^2 + y^2 + z^2) - \frac{1}{12}(x^4 + y^4 + z^4)$$
$$g = \nabla \Phi^\top \nabla \Phi$$

$$\tilde{\mathcal{Q}}_{xx} = \frac{9(z^2 - 1)(y^2 - 1)}{\det(g)(q(x) + q(y) + q(z) + 9)},$$

$$\tilde{\mathcal{Q}}_{yy} = \frac{9(z^2 - 1)(x^2 - 1)}{\det(g)(q(x) + q(y) + q(z) + 9)},$$

$$\tilde{\mathcal{Q}}_{zz} = \frac{9(x^2 - 1)(y^2 - 1)}{\det(g)(q(x) + q(y) + q(z) + 9)},$$


$$\tilde{\mathcal{Q}}_{xy} = \tilde{\mathcal{Q}}_{xz} = \tilde{\mathcal{Q}}_{yz} = 0,$$

$$q(x) = x^2(x^2 - 3)^2$$

Perturb mesh with uniform random noise to avoid possible super-convergence!


3D curvature

- Confirms theory for $k > 1$
- For $k = 0$ linear convergence is observed?!



3D curvature

- Confirms theory for $k > 1$
- For $k = 0$ linear convergence is observed?!
- Test only parts where theory indicates no convergence





3D curvature

$$F_1 : U \mapsto \frac{1}{2} \int_0^1 \sum_{E \in \mathring{\mathcal{E}}} \int_E \sigma_{\hat{\tau}_{\tilde{g}(t)} \hat{\tau}_{\tilde{g}(t)}} \Theta_E(\tilde{g}(t)) U_{\hat{\tau}_{\tilde{g}(t)} \hat{\tau}_{\tilde{g}(t)}} \omega_E(\tilde{g}(t)) dt$$

$$F_2 : U \mapsto -\frac{1}{2} \int_0^1 \sum_{E \in \mathring{\mathcal{E}}} \sum_{F \supset E} \int_E \sigma_{\hat{\tau}_{\tilde{g}(t)} \hat{\tau}_{\tilde{g}(t)}} [\![U_{\hat{\nu}_{\tilde{g}(t)} \hat{\mu}_{\tilde{g}(t)}}]\!]_F^E \omega_E(\tilde{g}(t)) dt$$

$$F_3 = F_1 + F_2$$



Summary & Outlook

- Definition of densitized distributional Riemann curvature tensor
- Analysis in the H^{-2} -norm via integral representation and Uhlenbeck trick
- Includes Gauss, scalar, and Ricci curvature tensor

Summary & Outlook

- Definition of densitized distributional Riemann curvature tensor
 - Analysis in the H^{-2} -norm via integral representation and Uhlenbeck trick
 - Includes Gauss, scalar, and Ricci curvature tensor
-
- Define appropriate FE to compute L^2 -representative and analyze in stronger norms
 - Investigate PDEs involving curvature fields, e.g. numerical relativity

Literature

-  LI: Regge Finite Elements with Applications in Solid Mechanics and Relativity, *PhD thesis, University of Minnesota* (2018).
-  BERCHENKO-KOGAN, GAWLIK: Finite element approximation of the Levi-Civita connection and its curvature in two dimensions, *Found Comput Math* (2022).
-  GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of curvature approximations via covariant curl and incompatibility for Regge metrics, *SMAI J. Comput. Math.* (2023).
-  GAWLIK, N.: Finite element approximation of scalar curvature in arbitrary dimension, *arXiv:2301.02159*.
-  GAWLIK, N.: Finite element approximation of the Einstein tensor, *arXiv:2310.18802*.
-  GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of distributional Riemann curvature tensor in any dimension, *arXiv:2311.01603*.

Literature

-  LI: Regge Finite Elements with Applications in Solid Mechanics and Relativity, *PhD thesis, University of Minnesota* (2018).
-  BERCHENKO-KOGAN, GAWLIK: Finite element approximation of the Levi-Civita connection and its curvature in two dimensions, *Found Comput Math* (2022).
-  GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of curvature approximations via covariant curl and incompatibility for Regge metrics, *SMAI J. Comput. Math.* (2023).
-  GAWLIK, N.: Finite element approximation of scalar curvature in arbitrary dimension, *arXiv:2301.02159*.
-  GAWLIK, N.: Finite element approximation of the Einstein tensor, *arXiv:2310.18802*.
-  GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of distributional Riemann curvature tensor in any dimension, *arXiv:2311.01603*.

Thank You for Your Attention!